



DECEMBER 11-12, 2025

۲۱ و ۲۰ آذر ماه ۱۴۰۴



دومین کنگره
پلازما پزشکی ایران

The 2nd Congress on Plasma Medicine

دبیرخانه دائمی کنگره
پلازما پزشکی ایران
www.plasmamedsym.ir



Plasma-Activated Media Induced Apoptosis in Monoculture and Co-Culture Systems of Breast Cancer Cells

Mahdiyeh Bakhtiyari Ramezani^{1*}, Marjan Mohamadali², and Ehsan Heidarian²

¹Plasma and Nuclear Fusion Research School, Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran

²Department of Biotechnology, Plasma Technology Development Company, Tehran, Iran

OPEN ACCESS

Citation:

Bakhtiyari Ramezani M, Mohamadali M, Heidarian E. Plasma-Activated Media Induced Apoptosis in Monoculture and Co-Culture Systems of Breast Cancer Cells. *Iran Biomed J. Supplementary* (2-2026): 11.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

ABSTRACT

Introduction: Breast cancer remains a major challenge due to the presence of aggressive and drug-resistant subtypes. Current therapeutic strategies have limited selectivity and severe side effects. Cold atmospheric plasma (CAP), which generates reactive oxygen species, induces apoptosis in cancer cells. In this study, the impact of CAP and plasma-activated media (PAM) on MDA-MB-231 and MCF-7 breast cancer cells, as well as their co-culture with human umbilical vein endothelial cells (HUVECs), was investigated.

Materials and Methods: CAP was generated using a plasma jet device operating with helium and argon gases under controlled conditions. The plasma treatment was applied in vitro to MCF-7 and MDA-MB-231 cells, along with HUVEC cells and their co-cultures. To assess cell viability, angiogenesis, and gene expression, we performed MTT, tube formation, and qRT-PCR. Apoptosis in treated cells was analyzed using Annexin V staining.

Results and Discussion: The results demonstrated significant efficacy of PAM in both monoculture and co-culture systems compared to direct plasma exposure ($p < 0.05$). Notably, PAM induced apoptosis and altered the expression of apoptosis-related genes, resulting in treated breast cancer cells.

Conclusion: Our findings suggest that PAM may offer a more effective and versatile approach for cancer treatment by a reduction in cell viability and modulating key biological processes such as apoptosis and angiogenesis.

Keywords: Breast cancer, CASPASE3, Cold atmospheric plasma, P53, Plasma-activated media, VEGF

Corresponding Author: Mahdiyeh Bakhtiyari Ramezani

Plasma and Nuclear Fusion Research School, Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran;

E-mail: mahdiyeh.bakhtiyari@gmail.com



Iranian Biomedical Journal Supplementary (February 2026): 11