



DECEMBER 11-12, 2025
۲۱ و ۲۰ آذر ماه ۱۴۰۴



دومین کنگره
پلازما پزشکی ایران

The 2nd Congress on Plasma Medicine

دبیرخانه دائمی کنگره
پلازما پزشکی ایران
www.plasmamedsym.ir



Effect of Cold Atmospheric Plasma on *Hsp90* Gene Expression in Terbinafine-Resistant Fungal Strains of the *Trichophyton mentagrophytes* Complex

Ameneh Shajari¹, Zahra Salehi², Mohammad Atyabi^{3*}

¹Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

²Department of Mycology, Pasteur Institute of Iran, Tehran, Iran

³Department of Nano Biotechnology, Pasteur Institute of Iran, Tehran, Iran

OPEN ACCESS

Citation:

Shajari A, Salehi Z, Atyabi M. Effect of Cold Atmospheric Plasma on *Hsp90* Gene Expression in Terbinafine-Resistant Fungal Strains of the *Trichophyton mentagrophytes* Complex. *Iran Biomed J. Supplementary* (2-2026): 45.

ABSTRACT

Introduction: The *Trichophyton mentagrophytes* complex, particularly terbinafine-resistant fungal strains, has emerged as a major cause of chronic and recurrent dermatophytosis, which is increasingly difficult to treat. These dermatophytes cause persistent infections of the skin, nails, and hair, often accompanied by inflammation, severe itching, keratin degradation, and reduced quality of life. In fungi, the molecular chaperone *Hsp90* is essential for stress tolerance, morphogenesis, and antifungal resistance. Cold atmospheric plasma (CAP), a non-thermal ionized gas, has recently gained attention for its notable antifungal activity. This study investigates the effect of CAP on *Hsp90* gene expression in terbinafine-resistant isolates of the *T. mentagrophytes* complex.

Materials and Methods: Fifty isolates of the *T. mentagrophytes* complex were tested for terbinafine susceptibility using broth microdilution, following the CLSI M38-A2 protocol. Isolates with minimum inhibitory concentrations above 0.25 µg/mL were classified as resistant. These resistant isolates, along with a standard sensitive strain (PFCC 5809), were exposed to a helium-based CAP jet, which was applied directly to fungal suspensions in 96-well microtiter plates. The expression levels of *Hsp90* before and after CAP exposure were quantified by real-time PCR using the $\Delta\Delta C_t$ method.

Results and Discussion: The baseline expression of *Hsp90* was 124.7-fold higher in resistant isolates compared to the sensitive strain. CAP treatment reduced *Hsp90* expression to 13.3-fold, which was significantly lower than baseline levels ($p < 0.005$).

Conclusion: CAP significantly downregulated *Hsp90* expression in terbinafine-resistant fungal strains. These findings suggest that CAP may serve as a promising adjunct antifungal strategy to enhance the management of resistant dermatophytosis.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

Keywords: Antifungal therapy, Cold atmospheric plasma, *Hsp90*, Terbinafine resistance, *Trichophyton mentagrophytes*

Corresponding Author: Mohammad Atyabi

Department of Nano Biotechnology, Pasteur Institute of Iran, Tehran, Iran; E-mail: mohammadatyabi@yahoo.com



Iranian Biomedical Journal Supplementary (February 2026): 45