



DECEMBER 11-12, 2025
۲۱ و ۲۰ آذر ماه ۱۴۰۴



دومین کنگره
پلاسما پزشکی ایران

The 2nd Congress on Plasma Medicine

دبیرخانه دائمی کنگره
پلاسما پزشکی ایران
www.plasmamedsym.ir



Selective Anticancer Effects of Cold Atmospheric Pressure Plasma Via RONS-Mediated Redox Dysregulation in Lung Cancer Cells

Mona Roshan^{1*}, Poopak Farnia², Hamid Reza Ghomi³, Saed Jafari⁴

¹Department of Physics, University Campus, University of Guilan, Rasht, Iran

²Mycobacteriology Research Centre, National Research Institute of Tuberculosis and Lung Disease, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Laser and Plasma Research Institute, Shahid Beheshti University, Tehran

⁴Department of Physics, Faculty of Science, University of Guilan, Rasht, Iran

OPEN ACCESS

Citation:

Roshan M, Farnia P, Ghomi HR, Jafari S. Selective Anticancer Effects of Cold Atmospheric Pressure Plasma Via RONS-Mediated Redox Dysregulation in Lung Cancer Cells. *Iran Biomed J. Supplementary* (2-2026): 46.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

ABSTRACT

Introduction: Cold atmospheric plasma (CAP) acts as a redox-modulating anticancer modality by generating reactive oxygen and nitrogen species (RONS). This process elevates lipid peroxidation (measured as malondialdehyde [MDA]), suppress superoxide dismutase (SOD) activity, and trigger oxidative stress-driven apoptosis. The integrated assessment of RONS, MDA, and SOD delineates the mechanistic basis of CAP-induced redox disequilibrium and cytotoxicity in lung cancer cells.

Materials and Methods: Human and murine lung cancer cells were exposed to helium-based CAP at voltages of 4 kV and 6 kV for 180 s using a flexible microplasma jet. Analyses were performed 24 h after the treatment. CAP-induced oxidative stress was analyzed by quantifying intracellular RONS, MDA, and SOD activity, while cytotoxicity was evaluated using the MTT assay.

Results and Discussion: CAP treatment markedly modulated the intracellular redox status in a voltage-dependent manner. At 6 kV, levels of intracellular RONS and MDA increased significantly, while SOD activity and cell viability (as measured by the MTT assay) decreased compared to 4 kV treatment. This outcome indicates that higher voltage led to greater disruption and induces apoptosis.

Conclusion: CAP treatment at 6 kV induced a marked rise in intracellular RONS and MDA levels, with a concurrent decline in SOD activity and cell viability, indicating a severe redox imbalance and apoptosis induction compared to 4 kV treatment.

Keywords: A549, Cold atmospheric pressure plasma, Helium, LL/2 cell lines, Lung cancer

Corresponding Author: Mona Roshan

Department of Physics, University Campus, University of Guilan, Rasht, Iran; E-mail: Mona_roshan_physics@yahoo.com



Iranian Biomedical Journal Supplementary (February 2026): 46